

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Conquest
Amine 625
Selective Herbicide

ACTIVE CONSTITUENT: 625 g/L 2,4-D PRESENT AS THE
DIMETHYLAMINE SALT

GROUP	I	HERBICIDE
-------	---	-----------

For the control of broadleaved weeds in fallow before direct drilling or sowing of cereals and pastures; and in cereal crops, pastures, sugar cane, peanuts and non-agricultural areas as per the Directions for Use.

**IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE USING THIS PRODUCT.**

Contents: 5 Litres
(20 L, 110 L, 200 L, 1000 L)

Conquest Agrochemicals Pty Ltd.
ABN 84 098 814 932
Suite 9, 41 Walters Drive
Herdsman Business Park
Osborne Park WA 6017

Telephone (08) 9244 1268
Facsimile: (08) 9446 4696

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Refillable containers : Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray, wear PVC or rubber apron, elbow length PVC gloves and face shield. When using the prepared spray, wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL
1 800 033 111
24 HOURS AUSTRALIA WIDE.

Conquest Agrochemicals Pty Ltd.

ABN 84 098 814 932
Suite 9, 41 Walters Drive
Herdsman Business Park
Osborne Park WA 6017

Telephone (08) 9244 1268
Facsimile: (08) 9446 4696

Batch No:
Date of Manufacture:
NRA Approval No: 55702/0602

CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Conquest Amine 625

Selective Herbicide

ACTIVE CONSTITUENT: 625 g/L 2,4-D PRESENT AS THE
DIMETHYLAMINE SALT

GROUP	I	HERBICIDE
-------	----------	-----------

For the control of broadleaved weeds in fallow before direct drilling or sowing of cereals and pastures; and in cereal crops, pastures, sugar cane, peanuts and non-agricultural areas as per the Directions for Use.

**IMPORTANT: READ THIS LEAFLET
BEFORE USING THIS PRODUCT.**

Conquest Agrochemicals Pty Ltd.

ABN 84 098 814 932
Suite 9, 41 Walters Drive
Herdsman Business Park
Osborne Park WA 6017

Telephone (08) 9244 1268
Facsimile: (08) 9446 4696

DIRECTIONS FOR USE

RESTRAINTS: DO NOT spray if rain seems likely within 6 hours or if strong winds prevail.
DO NOT apply crop or weeds are stressed due to dry or excessively moist conditions.

1. FIELD CROPS

SITUATION	WEEDS CONTROLLED	STATE	RATE VOL/HA	CRITICAL COMMENTS		
Wheat	Refer Weed Table	NSW, ACT, SA only	370 mL-1.7L	Apply after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller (NSW, SA only) Apply from tillering to boot stage (Vic only) Apply from mid tillering to before boot stage (Qld only) Apply at 5 leaf to fully tillered (Tas only)		
		Vic only	225 mL-1.4L			
		Qld only	560 mL-1.4L			
		Tas only	800 mL-1.4L			
		Barley	NSW, ACT SA only		320 mL-1.4L	
			Vic only		225 mL-1.4L	
			Qld only		560 mL-1.4L	
		Cereal Rye, Triticale	NSW, ACT SA only		Tas only	800 mL-1.4L
					NSW, ACT SA only	320 mL-1.4L
Oats	NSW, ACT, SA only	Vic only	225 mL-1.4L			
		Vic only	320 mL-800 mL			
Cereals: Wheat, Oats, Barley	Cape Tulip	WA only	640 mL-1.3L	Apply from the 5 leaf stage up to joining stage (Zadoks 15-33). Apply after the 6 leaf stage (Z.16) for Cranbrook, Jacup, Aroona and Spear Wheat and Mortlock Oats to avoid possible damage. DO NOT spray if lucerne is present.		
	Dock, Flatweed, Saffron Thistle		1.1L			
	Indian Hedge, Mustard London Rocket, Lupin, Matricaria, Rapistrum, Wild Radish		800 mL			
	Wild Turnip		640 mL			
	Capeweed, Doublegee, Erodium, London Rocket, Lupin, Mustard, Rapistrum, Wild Radish, Wild Turnip		200 mL/ha plus 500 mL/ha Flowable Diuron			
Wheat, Barley	Wild Radish	Vic, SA, NSW, ACT only	80 mL/ha plus 850g/ha Artfern Tribumax Herbicide	Spray 2-6 weeks after sowing and not later. DO NOT use on crops undersown with lucerne.		
Fallow, Stubble Spray prior to direct drilling or sowing a) Winter Cereals b) Winter Cereals, and Maize, Sweet corn, Peanuts	Refer Weed Table	Vic only	225 mL - 1.4L	Observe plant back periods given in the Table on this leaflet. Can be mixed with Conquest Chlorsulfuron 750 WG Herbicide, Paraquat or Spray Seed® where grasses are present. For Skeleton Weed, spraying should only be done 6-8 weeks before anticipated sowing date and subsequent cultivation limited to a minimum.		
		NSW, ACT only	560 mL - 1.7L			
		Qld only				
Maize, Sweetcorn	Refer to Weed Table	NSW, ACT, SA only	560 mL-1.1L	Apply when crop is 10-20 cm high and secondary roots are developing for an over the top spray. When crop is between 20 cm high and just before tasselling, spray with dropped nozzles to avoid chemical being sprayed into the whorl and on upper leaves		
		Qld only	560 mL-900 mL	Apply when crop is 10-30 cm high and secondary roots have developed. Use drop nozzles and direct the spray when crop exceeds 30 cm in height and before tasselling.		
		Tas only	800mL- 1.4 L	Apply when the crop is 15-30 cm high. Do not spray if the crop is showing signs of stress. Some leaf twisting may occur following application - crop recovers quickly.		

FIELD CROPS Continued.

Sorghum	Refer to Weed Table	NSW, ACT, SA, Vic only	560 mL-1.1L	Apply preferably when crop is at 3-6 fully expanded leaf stage but can be sprayed from 2-8 leaf stage. From 6 leaf stage onwards to within 2 weeks of flowering, crop can be sprayed with dropped nozzles to avoid chemical being sprayed into the whorl and on upper leaves.
		Qld only (except central Qld)	560-900mL	Apply when crop has 4 to 8 fully expanded leaves and secondary roots have developed.
Millet		NSW, ACT, SA, Vic only	560 mL-1.1L	Spray when secondary roots have been developed, when fully tillered and before heads start to form at the base of the tillers. Do not use on Panorama Millet or Panicum.
		Qld only	560-900mL	
Maize, Sweetcorn, Saccaline, Broom Millet, Millet,	Cape Tulip, Dock, Saffron Thistle, Indian Hedge Mustard, London Rocket, Lupin, Rapistrum, Wild Radish, Wild Turnip	WA only	1.1L	Spray when crop is 10-30 cm high and secondary roots have developed and before tasselling. Apply as direct spray to weeds.
Grain Sorghum				Apply when crop is 12 cm high. DO NOT apply between tassel and dough stage. Avoid spraying when in flower.
Sugar Cane (Q80, Q96, & H56 Varieties)	Bellvine	Qld, NSW, ACT only	280ml/100L water	Apply in spring using directed spray.
	Morning Glory		560ml-1.1L	Apply in summer using a high clearance tractor.
	Pink Convolvulus, Star of Bethlehem		1.1L	Apply in autumn by aircraft or misters.
Sugar Cane	Bindy Eye (Star Burr), Blue Top, Cobblers Pegs, Fleabanes, Jute, Leucas Needle Burr, Spear Thistle, Water Primrose, Ipomea Vines, Convolvulus Vines,	Qld only	1.8-3.5L	Add 170ml of Surfactant to 100L of spray mixture. Agitate well. DO NOT USE ON Q63, Q67, Q80 or Q96 VARIETIES.
	Chinese Mint, Blue Snakeweed		3.5L	
Peanuts	Broadleaf Weeds: except Noogoora Burr, Grasses: except Mossman Burr	Qld only	1.8L-3.6L	LOWER RATE: Apply as band spray as soon as possible after planting in a 55 cm band. HIGHER RATE: Apply as OVERALL SPRAY after planting and before crop emergence. Some crop damage may occur if heavy rain falls between application and crop emergence.
Harvest Aid or Salvage Spray - Winter Cereals - Maize & Sorghum	Desiccate Broadleaf Weeds Refer Weed Table	Qld, NSW & ACT only	1.2-1.7L	Apply after dough stage.
Bananas	To destroy Banana suckers	Qld only	160mL/10L water	Inject at a rate of 15mL per fully grown plant, 10mL per medium sized plant and 5mL for small suckers.
			320mL/100L water	Allow suckers from corms from treated plants to form broad adult leaves, then spray. Isolated spots may require a second spray.
Common Stylo forage or seed crops	Refer Weed Table		800mL	Apply post-emergence when weeds are 3 weeks old and crop is at least 3 weeks old.
Carribean Stylo forage or seed crops			800mL-1.6L	Apply post-emergence when crop is 3 weeks old.

2. PASTURES, NON AGRICULTURAL, RIGHTS OF WAY, INDUSTRIAL, LAWNS

SITUATION	WEEDS CONTROLLED	STATE	RATE VOL/HA	CRITICAL COMMENTS
Pastures and non-agricultural	Refer to Weed Table	NSW, ACT, QLD, SA, TAS only	560 mL-1.7L	Pasture legumes including lucerne, clovers and medics may be damaged unless well protected by grasses. Spot spraying is preferred.
	Galvanised Burr	NSW, ACT only	320mL/100L Water	Apply to young actively growing weeds. Ensure thorough and even coverage of plants. Note: Treated plants need to be burnt to destroy seeds.
	Amsinckia, Docks, Bindweed, Caltrop, Flatweed, Spear Thistle, Capeweed, Doublegee, Saffron, Thistle, Mustard, Wild Radish, Wild Turnip, Annual Thistles, Paterson's Curse, Heliotrope, Ragwort	WA only	1.1-2.4L	For pastures not containing legumes. Only seedling Docks, Spear Thistle and Saffron Thistle will be controlled. SUMMER WEEDS: Use low rate for seedlings, 1.6-2.4L/Ha for larger plants. Stock poisoning may occur when grazed after spraying if large amounts present, particularly Heliotrope. WINTER WEEDS: Use low rate for seedlings, 1.6-2.4L/Ha for larger plants. If stock present use spray/grazing rates.
	Afghan Melons		1.6L/ha plus 1% crop oil	Spray when plants are actively growing preferably before flowering or vining.
	Paddy Melons		800mL-1.1L	
	Prickly Saltwort (Roly Poly)		1.6L	Spray when plants are small.
	Stinkwort		1.6 - 3.2L + Surfactant	Best results are obtained when plants are small. Use high rate on larger plants.
	Dove Weed		3.2L	Spray after good emergence of seedlings.
Pastures, Rights of Way & Industrial	Boxthorn, Boneseed, Hawthorn	Vic, SA only	80 mL/10L water	Spot spraying. For Boneseed only, thoroughly wet plants and seedlings.
			Undiluted	Apply as an undiluted spray to freshly cut stumps.
	Groundsel	NSW, ACT, Qld, SA only	950ml/15L water	MISTING: Lightly wet plants.
			320 mL/100L water	HIGH VOLUME : Thoroughly wet plants
		Tas only	300ml/15L water	CUT STUMP: Swab the cut stump immediately. Apply by a pouring can or knapsack spray.
		NSW, ACT, Qld, SA, Tas only	2.9-4.4L	AERIAL APPLICATION : Spray when Groundsel is actively growing.
	Lantana	NSW, ACT, Qld, SA only	320 mL/100L water	Use a coarse spray with sufficient pressure to penetrate canopy and wet stems as well as foliage. Spray at the end of a wet summer (March to May). Defoliation should occur but respraying of new growth will be necessary in following Autumn. Broadcast grass seed and keep stock off following Summer to allow the pasture to establish. Damage may result to pasture legumes.
	Mother of Millions	NSW, ACT only	400 mL/100L water	Hand gun and Knapsack only. A thorough coverage of leaves and plantlets is necessary. Use Conquest Wetter 1000 at the rate of 1mL of surfactant per 1L of mixture.
Noogoora Burr, Weir Vine (Ipomoea), scarlet Pimpernel (seedlings only) White Eye (Mexican Clover)	Qld only	160 mL/100L water	In all cases apply to young, actively growing weeds, ensuring thorough coverage.	

PASTURE, NON AGRICULTURAL, RIGHTS OF WAY, INDUSTRIAL, LAWNS, Continued.

SITUATION	WEEDS CONTROLLED	STATE	RATE VOL/HA	CRITICAL COMMENTS
Pastures, Rights of Way & Industrial Cont.	Annual & Perennial Pigweed Artichoke Thistle, Bathurst Burr, Billygoat Weed, Blue Snake Weed, Burr Medic, Clockweed*, Fleabanes, Galvanised Burr, Hemlock, Hoary Cress ⁺ , Kyalinga Weed (Whisker Grass), Knobweed, Milky Cotton Bushes, Parthenium Weed, Paterson's Curse, Saffron Thistle, Star Burr, Thornapple, Variegated Thistle*	Qld only	320 mL/100L water	In all cases apply to young, actively growing weeds, ensuring thorough coverage. * Spray rosette stage. * Repeat spraying necessary.
	Rubber Vine		160 mL/10L water	Apply to freshly cut stump.
	Sesbania Pea		560mL-900mL	
	Water Hyacinth		3.5-5.3L	Apply in 2200 to 3300L water/ha
	Wild Tobacco Tree		240mL/15L water	Cut Stump Treatment: Swab cut stump within 1 hour of cutting. Apply by pouring can or knapsack sprayer.
Conservation Tillage - Direct Drilling, Surface Sowing or Fallow Maintenance	Charlock, Mustards, Shepherd's Purse, Saffron, Slender, Spear and Variegated Thistles*, Turnip Weed, Wild Radish, Wild Turnip	All States	560 mL-1.6L	Apply to actively growing young plants. Before sowing: Observe plant back periods given in the table on this leaflet. * Warning: Treated plants may become toxic to stock.
	Clover, Sorrel		1.1L/ha plus 700 mL-1L/ha of Banvel®	Apply to actively growing plants in autumn. Do not sow pasture seed for at least 30 days after application.
Pastures - Spray/Graze Techniques				PRECAUTION: An increased quantity of poisonous plants may be eaten by stock using Spray-Graze eg. Caltrop, Capeweed, Paterson's Curse and Variegated Thistle and deaths could result from causes such as nitrate poisoning. With Paterson's Curse, preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. Do not graze horses or pigs on Paterson's Curse.
	Amsinckia, Thistles, Capeweed, Doublegee, Mustard, Paterson's Curse, Wild Turnip, Wild Radish Docks, Geranium, Erodium	SA only	560 mL	Apply from 6 weeks after opening rains in Autumn until the end of August. Seven days after spraying stock paddock at 4-5 times normal rate, preferably with sheep. Maintain this level of grazing for 6 weeks or until pasture shows signs of over grazing, then return to normal stocking levels.
	Annual Thistles, Capeweed Doublegee, Mustards, Paterson's Curse, Wild Turnip, Saffron Thistle, Spear Thistle, Geranium, Slender Thistle	Tas, Vic only		Use high stocking rates in following Spring to prevent weeds from flowering. Repeat treatments may be required for 2-3 years for complete control.
	Amsinckia, Docks (seeding only), Capeweed, Doublegee, Mustard, Wild Radish, Wild Turnip, Paterson's Curse, Annual Thistles.	WA only		600-800 mL

PASTURE, NON AGRICULTURAL, RIGHTS OF WAY, INDUSTRIAL, LAWNS, Continued.

SITUATION	WEEDS CONTROLLED	STATE	RATE VOL/HA	CRITICAL COMMENTS
Pastures - Spray/Graze Techniques (cont)	Spear Thistle, Saffron Thistle	WA only	1.2L	Apply to Saffron Thistle at the end of September when plants are running up to flower. Sub-clovers may be damaged at this rate and use is not recommended for all medic pastures.
	Melons		1.6L + 1% oil	Heavy stocking on young plants sprayed with 800mL/ha provides effective control.
	Docks	Vic only	1.1L	Apply in September only and follow other recommendations above.
	Caltrop, Capeweed, Charlock, Mustards, Paterson's Curse, Shepherd's Purse, Saffron, Slender, Spear or Variegated Thistle*, Turnip Weed, Wild Radish, Wild Turnip	NSW, ACT only	280 mL-1.1L	Spray actively growing 6-8 week old weeds. Introduce stock 7-10 days after spraying, preferably sheep (cattle are less effective). Stocking rate should be at least 5 times heavier than normal until weeds have been reduced, but before survival of desirable pasture species is threatened. Lucerne and Medics may be damaged and should be grazed short before spraying. Other legumes may be affected.
Lawns, Playing fields	Refer Weed Table	Qld, NSW only	1.5-3 mL/1L water or 1.6-3.2L/ha	Wet foliage thoroughly. DO NOT mow lawn for 1 week before and at least 1 week after application. DO NOT use on Buffalo grass (WA only).
		WA only	40 mL/10-15L water/100 m ²	

3. SPOT SPRAYING

SITUATION	WEEDS CONTROLLED	STATE	MIXING RATES/CRITICAL COMMENTS
High volume spraying	Refer to Weed Table	All States	Add 1/10th of rate on Weed Table to 150 litres of water. Each 150 litres of mix will cover 1000m ² (1/10th ha) eg. if rate in Weed Table is 1.4L, use 140 mL/150L water.
Knapsack Application	Refer to Weed Table	All States	Add 1/100th of rate on Weed Table to 10 litres of water. Each 10 litres of mix will cover 100m ² (1/100th ha) eg. if rate in Weed Table is 1.4L, use 14 mL/10L water.

MAXIMUM TOLERANCE FOR USE IN WHEAT, BARLEY, OATS, AND TRITICALE INCLUDING TOLERANCE FOR UNDERSOWN LEGUMES							
Crop	Qld	NSW, ACT	Vic	Vic	SA	WA	Tas
			<u>Early Tillering</u>	<u>Tillered to Boot Stage</u>		<u>Tillered to Boot (Z15-35)</u>	<u>Tillered to Boot (Z15-35)</u>
Cereal Rye					1.4L/ha		
Wheat	1.8L/ha	1.7L/ha	225mL/ha	1.4L/ha	1.7L/ha	1.7L/ha	1.7L/ha
Barley	1.4L/ha	1.4L/ha	225mL/ha	1.4L/ha	1.4L/ha	1.4L/ha	1.4L/ha
Oats	-	800mL/ha	225mL/ha	800mL/ha	800mL/ha	1.3L/ha	-
Triticale	-	1.4L/ha	225mL/ha	-	1.4L/ha	1.7L/ha	800mL/ha
Undersown clovers	-	-	225mL/ha	-	-	680mL/ha	560mL/ha
Undersown Medics	-	-	225mL/ha	-	-	NIL	560mL/ha
Undersown Lucerne	-	-	225mL/ha	-	-	NIL	-

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THE LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD:

GRAZING: PASTURE AND CEREAL CROPS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION

CROP HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

In Tasmania, this product may only be used from 15 April to 15 September unless otherwise permitted by the registrar of pesticides.

WEED TABLE

Where weeds are to be sprayed in a CROP or PASTURE, use only the rate given for the particular crop or situation indicated under the Directions For Use table. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed. This product should only be used in those States where a rate or range of rates is indicated for the particular weeds listed. The rates listed in the weed table below are for use where weeds are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage.

This product should only be used in those states where a rate or range of rates is indicated for the particular weeds listed.

WEEDS	APPLICATION RATE PER HECTARE							CRITICAL COMMENTS
	CROP						PASTURE	
	VIC	NSW, ACT	SA	QLD	TAS	WA	NSW, SA, QLD, TAS ONLY	
Amaranthus spp.	-	560mL-1.1L	-	900mL	-	-	-	Spray young plants.
Amsinckia	-	-	-	-	-	1.1L	-	
Apple of Peru	-	560mL - 1.1L	-	900mL	-	-	-	Spray young plants. Susceptible when young.
Bathurst Burr	-	800mL-1.1L	1.7 - 2.2L	900mL	-	-	800mL - 1.1L Not SA	Spray seedlings only.
Bellvine	-	1.8L	-	1.8L	-	-	-	Spray before seeding. Advanced stages susceptible
Billygoat Weed	-	3.4L	-	3.4L	-	-	-	Spray at young stage.
Bindweed	-	-	-	-	-	1.1L	-	
Blackberry Nightshade	-	560mL - 1.1L	-	900mL	-	-	-	
Blackeyed Susan	-	1.8L	-	1.8L	-	-	-	Apply at pre-flowering, preferably young stages.
Blue Snakeweed	-	1.8L	-	1.8L	-	-	-	Spray seedlings at young stages only.
California Burr	-	800mL-1.1L	-	900mL	-	-	800mL-1.1L not SA	Spray seedlings only.
Cape Tulip	-	-	-	-	-	640mL -1.3L	-	Low rate for cormils only*.
Capeweed	1.4L	-	1.7L	-	1.4L	1.1L	1.7-2.8L	Spray seedlings to rosette stage.
Caltrop	-	800mL-1.7L	-	900mL	-	1.1L	-	Moderately susceptible
Caster Oil Plant	-	3.4L	-	3.4L	-	-	-	Spray at young stage.
Charlock	560mL-800mL	560mL-1.1L	560mL	-	1.4L	-	800mL-1.1L	Spray at rosette stage
Clover	-	1.2L	-	-	-	-	-	
Cobbler's Pegs	-	1.8L	-	1.8L	-	-	-	Apply at pre-flowering, preferably young stages.
Common Ice Plant	-	-	1.1L	-	-	-	-	
Common Sida	-	1.8L	-	1.8L	-	-	-	Spray seedlings or young stages only.
Common Sowthistle	-	1.8L	-	1.8L	1.4L	-	-	Apply at pre-flowering, preferably young stages.
Docks	1.4	-	1.4L	1.4L	1.4L	1.1L	3.2L SA only	Spray at multiple leaf stage - effective only on seedlings.
Doveweed	-	-	-	-	-	1.1L	-	
Fat Hen	-	560mL-1.7L	-	900mL	1.4L	-	-	Spray pre-flowering.
Flannel weed	-	1.8L	-	1.8L	-	-	-	Spray seedlings or young stages only.
Flat Weed	-	-	-	-	-	1.1L	-	
Fumitory – red	-	-	1.7L	-	-	-	-	
Fumitory – white	800mL	-	560mL	-	-	-	-	Spray at multiple leaf stage.
Heliotrope	-	-	-	-	-	1.1L	-	
Hexham Scent or Melilotus	1.4L	-	1.1L	1.4L	-	-	1.1-1.7L	Spray multiple leaf stage before seeding.
Hoary Cress	900mL-1.4L	1.1-1.7L	1.4L	1.4L	-	-	1.4-1.7L	Spray rosettes at pre-flowering.
Hogweed/ Wireweed	1.4L	-	-	1.4L	-	-	-	Spray at multiple leaf stage (Vic) Spray at seedling and young plant stage (Qld).

WEEDS	APPLICATION RATE PER HECTARE							CRITICAL COMMENTS
	CROP						PASTURE	
	VIC	NSW, ACT	SA	QLD	TAS	WA	NSW, SA, QLD, TAS ONLY	
Horehound	-	-	1.4L	-	-	-	2.2-3.2L	Spray seedlings.
Indian Hedge Mustard	-	-	-	-	1.4L	1.1L	-	
Khaki Weed	-	-	-	-	-	-	1.1-2.2L not SA	Spray seedlings only
Knobweed	-	-	-	1.8-3.4L	-	-	-	Lower rate for seedlings; higher rate for later stages.
Lincoln Weed	-	-	1.7L	-	-	-	-	Spray early rosettes
London Rocket	-	-	-	-	-	1.1L	-	
Lupins	-	800mL-1.7L	-	-	-	1.1L	-	
Melons – Camel, Paddy	-	560mL-1.1L	-	-	-	-	-	
Mexican Poppy	-	-	-	1.4L	-	-	-	Spray seedlings - plants become more resistant with age.
Mintweed	-	1.1L	-	900mL	-	-	-	Spray seedlings - resistant in later stages.
Morning Glory	-	1.8L	-	1.8L	-	-	-	Spray at seedling to flowering stage.
Mustards	225-560mL	560ml-1.1L	560ml-1.4L	900mL	-	800mL	560ml-1.1L	Spray at 2-4 leaf up to rosette stage.
Needle Burr	-	1.8L	-	1.8L	-	-	-	Apply at pre-flowering preferably young stages.
New Zealand Spinach	-	1.1-1.7L	-	-	-	-	-	
Noogoora Burr	-	800mL-1.1L	-	900mL	-	-	800mL-1.1L not SA	Spray seedlings only
Paterson's Curse	-	1.1-1.7L	-	1.4L	-	1.3L	1.7-2.2L	Spray rosettes or before plants have 10 leaves. Later stages harder to kill.
Pinkburr (Pink Flowered Burr)	-	1.8L	-	1.8L	-	-	-	Spray seedlings or young stages only.
Potato Weed	-	560mL-1.1L	-	900mL	-	-	-	
Purpletop	-	3.4L	-	3.4L	-	-	-	Spray at young stage.
Radish	-	-	-	-	-	1.1L	-	
Ragwort	-	-	-	-	3.8L	1.1L	-	Spray up to early rosette stage (Tas only).
Rapeseed	-	800mL-1.7L	-	-	-	-	-	
Rapistrum	-	-	-	-	-	1.1L	-	
Rough Poppy	-	1.1L	-	-	-	-	-	
Safflower	-	560mL-1.1L	-	-	-	-	-	
Shepherds Purse	-	1.1-1.7L	-	-	1.4L	-	800mL-1.1L	Spray young rosettes
Siratro (Purple Bean)	-	1.8L	-	1.8L	-	-	-	Spray seedlings at young stages only.
Skeleton Weed	1.4L	1.1-1.7L	1.4L	-	-	-	-	Spray rosettes before aerial growth commences.
Sorrel	1.4L	1.7L	1.4L	-	-	-	-	Only moderately susceptible
Speedwell - Ivy leaf	-	-	1.1L	-	-	-	-	
Spinyhead Sida	-	1.8L	-	1.8L	-	-	-	Spray seedling or young stages only.
Starburr	-	1.8L	-	1.8L	-	-	-	Spray before seeding, advanced stages susceptible.
Spiny Emex	-	-	-	1.4L	-	-	-	Only young plants are susceptible.
Star of Bethlehem (Cupid's Flower)	-	-	-	1.8L	-	-	-	Spray before seeding, advanced stages susceptible.
Stinkwort	-	800mL-1.4L	-	-	-	-	-	
Storksbill/Erodium	-	-	-	-	1.4L	-	1.6-3.2L	Spray seedlings to young rosettes.
Sunflower (seedlings)	1.4L	560mL-1.4L	-	900mL	-	-	-	

WEEDS	APPLICATION RATE PER HECTARE							CRITICAL COMMENTS
	CROP						PASTURE	
	VIC	NSW, ACT	SA	QLD	TAS	WA	NSW, SA, QLD, TAS ONLY	
Thistles - Annual	-	-	-	-	-	1.1L	-	
- Californian	-	-	-	-	2.7L	-	3.2-3.8L	Repeated applications may be necessary (NSW, Tas only).
- Saffron	1.1L	560mL-1.7L	1.4L	1.4L	950mL	1.1L	1.1-1.7L	Low rate only sufficient to control weeds in crops at rosette stage when sprayed early.
- Slender/Shore	-	800mL-1.7L	-	-	1.4L	-	1.1L	Suppression only.
- Soldier	1.4L	-	-	-	-	-	1.1-1.6L not NSW, Tas	Spray young rosettes
- Spear	560mL	-	-	-	1.4L	-	1.1-1.6L	Spray young rosettes
- Star	-	-	-	-	-	-	1.6-3.2L SA only	Use higher rate as flower stalk appears.
- Variegated	-	560mL-1.7L	-	900mL	1.4L	-	1.1-1.7L	Spray at rosette stage.
Thornapple	-	800mL-1.1L	-	-	-	-	1.6-2.4 not SA	Spray seedlings only.
Tridax (Tridax Daisy)	-	1.8L	-	1.8L	-	-	-	Spray seedling or young stages only.
Turnip Weed/Rapistrum	-	560mL-1.1L	-	560mL	-	800mL	560ml-1.1L	
Vetches/Tares	1.4L	-	1.1L	-	-	-	-	Spray at multiple leaf stage.
Wards Weed	-	-	1.1L	-	-	-	-	
Wild Cabbage	1.4L	-	-	-	-	-	-	Spray multiple leaves.
Wild Poppy	560mL	-	-	-	-	-	1.1-1.7L	Spray rosettes.
Wild Radish	1.4L	1.4-1.7L	1.4L	900mL	1.4L	800mL	800mL-1.1L	Spray up to young rosette stage.
Wild Turnip	225-560mL	560mL-1.1L	320mL	-	1.4L	640ml	560mL-1.1L	Spray 2-4 leaf up to rosette stage.

PLANT BACK DAYS FOR Conquest Amine 625 Selective Herbicide

CROP	RATES		
	Up to 560 mL/ha	560 mL - 1.1L/ha	1.1L - 1.7L/ha
Balansa Clover	7	7	10
Barley %	1	1	3
Chickpeas #	7	14	21
Cotton	10	14	21
Faba Beans	7	7	10
Field Peas	7	14	14
Lentils	7	7	10
Linseed	7	7	14
Lucerne	7	7	10
Lupins +	7	14	21
Medics	7	7	10
Narbon Beans	7	7	10
Navy Bean	10	10	14
Oats	3	3	7
Perennial Ryegrass	7	7	10
Persian Clover	7	7	10
Phalaris	7	7	10
Canola/Rapeseed #	14	21	28
Rice	7	7	14
Safflower #	7	14	21
Sorghum @	3	7	10
Soybean	14	14	21
Sub-Clover	7	7	10
Sunflower @	7	10	14
Triticale %	1	3	7
Vetch	7	7	10
Wheat %	1	3	7
White Clover	7	7	10

IMPORTANT - WHEN APPLIED TO DRY SOILS, AT LEAST 15 MM (1/2 INCH) OF RAIN MUST FALL PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.

NOTES:

% In Queensland, no rainfall is required to fall prior to commencement of Plant Back Period for Wheat, Barley and Triticale.

In Queensland, planting of Canola/Rapeseed, Chickpeas and Safflower must be delayed for at least 14 days following rainfall of at least 15 mm

@ In Central Queensland and when using 800mL/ha or less of Conquest Amine 625 Selective Herbicide, the plant back period for Sorghum and Sunflower is 1 day irrespective of rainfall.

+ In WA the plant back period for Lupins at all rates is 28 days.

GENERAL INSTRUCTIONS:

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions.

Resistant Weeds Warning:

GROUP	I	HERBICIDE
-------	---	-----------

Conquest Amine 625 Selective Herbicide is a member of the Phenoxy group of herbicides. It has the disrupters of plant cell growth mode of action. For weed resistance management Conquest Amine 625 Selective Herbicide is a group I Herbicide.

Some naturally-occurring weed biotypes resistant to Conquest Amine 625 Selective Herbicide and other disrupters of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Conquest Amine 625 Selective Herbicide or other disrupters of plant cell growth herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Conquest Agrochemicals Pty Ltd accepts no liability for any losses that may result from the failure of Conquest Amine 625 Selective Herbicide to control resistant weeds.

Application Information:

BOOM SPRAYING – Use 30–120L/ha of water.

AERIAL SPRAYING – Use 10–90L/ha of water.

Equipment Maintenance and Usage:

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants, unless it has been well washed out with hot soapy water or 1% solution of ammonia, followed by several clear water rinses.

Compatibility:

This product can be tank mixed with dicamba, Flowable Diuron, Conquest Chlorsulfuron 750 WG Herbicide, Paraquat, Conquest Metsulfuron 600 WG Herbicide, Atrazine, Sprayseed®, Tordon 50–D® and Methabenzthiazuron.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS:

DO NOT spray in high winds. DO NOT spray cereals if lucerne is present. DO NOT spray crops or weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result.

Drift Warning: DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants /crops, cropping lands or pastures.

Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

PROTECTION OF LIVESTOCK:

Low hazard to bees. May be applied at any time as recommended in the Directions for Use

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Refillable containers : Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray, wear PVC or rubber apron, elbow length PVC gloves and face shield. When using the prepared spray, wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

For specialist advice in an emergency dial
1 800 033 111
24 hours Australia wide.

Conquest Agrochemicals Pty Ltd.

ABN 84 098 814 932
Suite 9, 41 Walters Drive
Herdsman Business Park
Osborne Park WA 6017

Telephone (08) 9244 1268
Facsimile: (08) 9446 4696

NRA Approval No: 55702/0602